

## 70: Discrete Math and Probability Theory

Programming + Microprocessors  $\equiv$  Superpower!

What are your super powerful programs/processors doing?

Logic and Proofs!

Induction  $\equiv$  Recursion.

What can computers do?

Work with discrete objects.

**Discrete Math**  $\implies$  immense application.

Computers learn and interact with the world?

E.g. machine learning, data analysis, robotics, ...

**Probability!**

See note 1, for more discussion.

## Admin

Course Webpage: <http://www.eecs70.org/>

Explains policies, has office hours, homework, midterm dates, etc.

Two midterms, final.

midterm 1 before drop date.

midterm 2 before grade option change.

Questions/Announcements  $\implies$  piazza:

[piazza.com/berkeley/fall2016/cs70](http://piazza.com/berkeley/fall2016/cs70)

## Instructors

Instructor: Sanjit Seshia.

Professor of EECS (office: 566 Cory)

Starting 12th year at Berkeley.

PhD: in Computer Science, from Carnegie Mellon University.

Research: Formal Methods (a.k.a. Computational Proof Methods)

applied to cyber-physical systems (e.g. "self-driving" cars), computer security, ...

Taught: 149, 172, 144/244, 219C, EECS149.1x on edX, ...

## CS70: Lecture 1. Outline.

Today: Note 1. (Note 0 is background. Do read/skim it.)

The language of proofs!

1. Propositions.
2. Propositional Forms.
3. Implication.
4. Truth Tables
5. Quantifiers
6. More De Morgan's Laws

## Instructors

Jean Walrand – Prof. of EECS – UCB  
257 Cory Hall – walrand@berkeley.edu



I was born in **Belgium**<sup>(1)</sup> and came to Berkeley for my PhD. I have been teaching at UCB since 1982.

My wife and I live in Berkeley. We have two daughters (UC alumni – Go Bears!). We like to ski and play tennis (both poorly). We enjoy classical music and jazz.

My research interests include stochastic systems, networks and game theory.

(1)



## Propositions: Statements that are true or false.

$\sqrt{2}$  is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Jon Stewart is a good comedian

All evens  $> 2$  are unique sums of 2 primes

$4 + 5$

$x + x$

**Proposition** True

**Proposition** True

**Proposition** False

**Proposition** False

**Not a Proposition**

**Proposition** False

**Not a Proposition.**

**Not a Proposition.**

Again: "value" of a proposition is ... True or False

## Propositional Forms.

Put propositions together to make another...

Conjunction ("and"):  $P \wedge Q$

" $P \wedge Q$ " is True when both  $P$  and  $Q$  are True . Else False .

Disjunction ("or"):  $P \vee Q$

" $P \vee Q$ " is True when at least one  $P$  or  $Q$  is True . Else False .

Negation ("not"):  $\neg P$

" $\neg P$ " is True when  $P$  is False . Else False .

Examples:

$\neg$  "(2+2 = 4)" - a proposition that is ... False

"2+2 = 3"  $\wedge$  "2+2 = 4" - a proposition that is ... False

"2+2 = 3"  $\vee$  "2+2 = 4" - a proposition that is ... True

## Propositional Forms: quick check!

$P = \sqrt{2}$  is rational"

$Q =$  "826th digit of pi is 2"

$P$  is ...False .

$Q$  is ...True .

$P \wedge Q$  ... False

$P \vee Q$  ... True

$\neg P$  ... True

## Put them together..

Propositions:

$P_1$  - Person 1 rides the bus.

$P_2$  - Person 2 rides the bus.

....

Suppose we can't have either of the following happen; That either person 1 or person 2 ride the bus and person 3 or 4 ride the bus. Or that person 2 or person 3 ride the bus and that either person 4 ride the bus or person 5 doesn't.

Propositional Form:

$\neg(((P_1 \vee P_2) \wedge (P_3 \vee P_4)) \vee ((P_2 \vee P_3) \wedge (P_4 \vee \neg P_5)))$

Who can ride the bus?

What combinations of people can ride the bus?

This seems ...complicated.

We need a way to keep track!

## Truth Tables for Propositional Forms.

$P$	$Q$	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

$P$	$Q$	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

One use for truth tables: Logical Equivalence of propositional forms!

Example:  $\neg(P \wedge Q)$  logically equivalent to  $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

$P$	$Q$	$\neg(P \wedge Q)$	$\neg P \vee \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

DeMorgan's Law's for Negation: distribute and flip!

$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$        $\neg(P \vee Q) \equiv \neg P \wedge \neg Q$

## Implication.

$P \implies Q$  interpreted as

If  $P$ , then  $Q$ .

True Statements:  $P, P \implies Q$ .

Conclude:  $Q$  is true.

Example: Statement: If you stand in the rain, then you'll get wet.

$P =$  "you stand in the rain"

$Q =$  "you will get wet"

Statement: "Stand in the rain"

Can conclude: "you'll get wet."

## Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is False if  $P$  is True and  $Q$  is False .

False implies nothing

$P$  False means  $Q$  can be True or False

Anything implies true.

$P$  can be True or False when  $Q$  is True

If chemical plant pollutes river, fish die.

If fish die, did chemical plant polluted river?

Not necessarily.

$P \implies Q$  and  $Q$  are True does not mean  $P$  is True

Instead we have:

$P \implies Q$  and  $P$  are True does mean  $Q$  is True .

Be careful out there!

Some Fun: use propositional formulas to describe implication?

$((P \implies Q) \wedge P) \implies Q$ .

## Implication and English.

$$P \implies Q$$

- ▶ If  $P$ , then  $Q$ .
- ▶  $Q$  if  $P$ .
- ▶  $P$  only if  $Q$ .
- ▶  $P$  is sufficient for  $Q$ .
- ▶  $Q$  is necessary for  $P$ .

## Variables.

Propositions?

- ▶  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$ .
- ▶  $x > 2$
- ▶  $n$  is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g.,  $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- ▶  $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- ▶  $R(x) = "x > 2"$
- ▶  $G(n) = "n \text{ is even and the sum of two primes}"$

Next: Statements about boolean valued functions!!

## Truth Table: implication.

$P$	$Q$	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

$P$	$Q$	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

$$\neg P \vee Q \equiv P \implies Q.$$

These two propositional forms are logically equivalent!

## Quantifiers..

**There exists quantifier:**

$(\exists x \in S)(P(x))$  means " $P(x)$  is true for some  $x$  in  $S$ "

**Wait! What is  $S$ ?**

$S$  is the **universe**: "the type of  $x$ ".

Universe examples include..

- ▶  $N = \{0, 1, \dots\}$  (natural numbers).
- ▶  $Z = \{\dots, -1, 0, 1, \dots\}$  (integers)
- ▶  $Z^+$  (positive integers)
- ▶ See note 0 for more!

## Contrapositive, Converse

- ▶ Contrapositive of  $P \implies Q$  is  $\neg Q \implies \neg P$ .
  - ▶ If the plant pollutes, fish die.
  - ▶ If the fish don't die, the plant does not pollute. (contrapositive)
  - ▶ If you stand in the rain, you get wet.
  - ▶ If you did not stand in the rain, you did not get wet. (not contrapositive!) converse!
  - ▶ If you did not get wet, you did not stand in the rain. (contrapositive.)

Logically equivalent! Notation:  $\equiv$ .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- ▶ Converse of  $P \implies Q$  is  $Q \implies P$ .  
If fish die the plant pollutes.  
Not logically equivalent!
- ▶ **Definition:** If  $P \implies Q$  and  $Q \implies P$  is  $P$  if and only if  $Q$  or  $P \iff Q$ .  
(Logically Equivalent:  $\iff$  .)

## Quantifiers..

**There exists quantifier:**

$(\exists x \in S)(P(x))$  means " $P(x)$  is true for some  $x$  in  $S$ "

For example:

$$(\exists x \in N)(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

**Much shorter to use a quantifier!**

**For all quantifier;**

$(\forall x \in S)(P(x))$ . means "For all  $x$  in  $S$   $P(x)$  is **True**."

Examples:

"Adding 1 makes a bigger number."

$$(\forall x \in N)(x + 1 > x)$$

"the square of a number is always non-negative"

$$(\forall x \in N)(x^2 \geq 0)$$

## Quantifiers are not commutative.

- ▶ Consider this English statement: "there is a natural number that is the square of every natural number", i.e the square of every natural number is the same number!

$$(\exists y \in \mathbb{N})(\forall x \in \mathbb{N})(y = x^2) \quad \text{False}$$

- ▶ Consider this one: "the square of every natural number is a natural number"...

$$(\forall x \in \mathbb{N})(\exists y \in \mathbb{N})(y = x^2) \quad \text{True}$$

## Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

By DeMorgan's law,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

English: there is an  $x$  in  $S$  where  $P(x)$  does not hold.

**What we do in this course! We consider claims.**

**Claim:**  $(\forall x) P(x)$  "For all inputs  $x$  the program works."

For **False**, find  $x$ , where  $\neg P(x)$ .

Counterexample.

Bad input.

Case that illustrates bug.

For **True**: prove claim. Next lectures...

## Negation of exists.

Consider

$$\neg(\exists x \in S)(P(x))$$

Equivalent to:

$$\neg(\exists x \in S)(P(x)) \iff \forall(x \in S)\neg P(x).$$

English: means that for all  $x$  in  $S$ ,  $P(x)$  does not hold.

## Which Theorem?

Theorem:  $\forall n \in \mathbb{N} (n \geq 3 \implies \neg(\exists a, b, c \in \mathbb{N} a^n + b^n = c^n))$

Which Theorem?

Fermat's Last Theorem!

Remember Right-Angled Triangles: for  $n = 2$ , we have 3,4,5 and 5,7, 12 and ... (Pythagorean triples)

1637: Proof doesn't fit in the margins.

1993: Wiles ...(based in part on Ribet's Theorem)

DeMorgan Restatement:

Theorem:  $\neg(\exists n \in \mathbb{N} \exists a, b, c \in \mathbb{N} (n \geq 3 \wedge a^n + b^n = c^n))$

## Summary.

Propositions are statements that are true or false.

Propositional forms use  $\wedge, \vee, \neg$ .

The meaning of a propositional form is given by its truth table.

Logical equivalence of forms means same truth tables.

Implication:  $P \implies Q \iff \neg P \vee Q$ .

Contrapositive:  $\neg Q \implies \neg P$

Converse:  $Q \implies P$

Predicates: Statements with "free" variables.

Quantifiers:  $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: "Flip and Distribute negation"

$$\neg(P \vee Q) \iff (\neg P \wedge \neg Q)$$

$$\neg \forall x P(x) \iff \exists x \neg P(x).$$

Next Time: proofs!